



Simon Bangiyev, DDS, MD

Diplomate of the American Board of Oral and Maxillofacial Surgery

**CONSENT FOR ORAL SURGERY TREATMENT IN PATIENTS WHO HAVE RECEIVED INTRAVENOUS BIPHOSPHONATE DRUGS, ANTIRESORPTIVE DRUGS, OR ANTIANGIOGENIC DRUGS**

\_\_\_\_\_  
Patient's Name

\_\_\_\_\_  
Date of Birth

**Please initial each paragraph after reading. If you have any questions, please ask your doctor BEFORE initialing.**

\_\_\_ 1. I understand I have been treated with medication to help treat my specific cancer diagnosis:

- IV Bisphosphonate Drugs (Zometa/Aredia)
- Antiresorptive Drugs (Denosumab)
- Antiangiogenic Drugs

I have been informed that there is a risk of developing a complication called osteonecrosis of the bone (bone cell death) that can occur with oral surgical treatment including routine tooth extraction. Jaw bones usually heal completely. In some patients taking these medications, the ability of the bone to heal may be altered, interfering with the jaw's ability to heal normally. The research shows this risk occurs in less than 3% of patients having a tooth extracted. Any invasive procedure affecting the bone can increase the risk of osteonecrosis, so it is important to understand these risks before proceeding with any surgical procedure.

\_\_\_ 2. I understand that the risk of osteonecrosis can be increased by certain medical conditions including diabetes, immune suppression, cancer, as well as social habits like chronic smoking.

\_\_\_ 3. My medical/dental history is very important. I have provided you with a complete list of medications that I have received or taken before, and are receiving or taking now. I understand the importance of providing my doctor with a thorough and accurate medical history, including names of physicians I am seeing now or have seen in the past.

\_\_\_ 4. The decision to stop IV Bisphosphonate drug therapy before dental treatment will not lessen the risk of developing Osteonecrosis. Stopping the anti-angiogenic medications prior to dental treatment may improve healing and should be reviewed with your treating doctors.

\_\_\_ 5. Antibiotic therapy may be used to help control possible post-operative infection. For some patients, taking antibiotics may cause allergic responses or have unwanted side effects such as stomach discomfort, diarrhea, swelling of the colon, etc.

\_\_\_ 6. Even with all the precautions we take, there may be delayed healing, necrosis of the jaw bone, loss of bone and soft tissues, infection, fracture of the jaw due to a medical condition, oral-cutaneous fistula (open draining wounds), or other significant complications.

Patient's Initials \_\_\_\_\_

